



Contact CR-Warped Product Submanifolds of Kenmotsu Manifolds

V.A. Khan, K.A. Khan and Sirajuddin

Abstract : Recently B.Y. Chen [4] studied warped products which are CR-submanifolds in Kaehler manifolds. B. Sahin [8] extended the above result for warped product semi-slant submanifolds $N_\theta \times_f N_T$ of a Kaehler manifold. In the present paper we have investigated the existence of doubly as well as CR-warped product submanifolds of Kenmotsu manifolds.

Keywords : Doubly warped product, warped product, contact CR-submanifold, Kenmotsu manifold.

2000 Mathematics Subject Classification : 53C40; 53B25

1. Preliminaries

A $(2m + 1)$ -dimensional Riemannian manifolds (\bar{M}, g) is said to be a *Kenmotsu manifold* if it admits an endomorphism ϕ of its tangent bundle $T\bar{M}$, a vector field ξ and a 1-form η satisfying:

$$\left. \begin{aligned} \phi^2 X &= -X + \eta(X)\xi, \quad \eta(\xi) = 1, \quad \phi(\xi) = 0, \quad \eta \circ \phi = 0, \\ g(\phi X, \phi Y) &= g(X, Y) - \eta(X)\eta(Y), \quad \eta(X) = g(X, \xi), \\ (\bar{\nabla}_X \phi)Y &= g(\phi X, Y)\xi - \eta(Y)\phi X, \quad \bar{\nabla}_X \xi = X - \eta(X)\xi \end{aligned} \right\} \quad (1.1)$$

for any vector fields X, Y on \bar{M} , where $\bar{\nabla}$ denotes the Riemannian connection with respect to g .

Now, let M be a submanifold immersed in \bar{M} . We also denote by g the induced metric on M . Let TM be the Lie algebra of vector fields in

M and $T^\perp M$ the set of all vector fields normal to M . Denote by ∇ the Levi-Civita connection on M . Then Gauss-Weingarten formulas are given by

$$\bar{\nabla}_X Y = \nabla_X Y + h(X, Y) \quad (1.2)$$

$$\bar{\nabla}_X N = -A_N X + \nabla_X^\perp N \quad (1.3)$$

for any $X, Y \in TM$ and any $N \in T^\perp M$, where ∇^\perp is the connection in the normal bundle, h is the second fundamental form of M and A_N is the Weingarten endomorphism associated with N . The second fundamental form h and the shape operator A are related by

$$g(A_N X, Y) = g(h(X, Y), N) \quad (1.4)$$

A submanifolds M tangent to ξ is called *contact CR-submanifold* if it admits an invariant distribution D whose orthogonal complementary distribution D^\perp is anti-invariant i.e., $TM = D \oplus D^\perp \oplus \langle \xi \rangle$ with $\phi(D_p) \subseteq D_p$ and $\phi(D_p^\perp) \subset T_p^\perp M$, for every $p \in M$. It is clear that contact CR-submanifold is a special case of semi-slant submanifolds.

For any $X \in TM$, we write

$$\phi X = PX + FX \quad (1.5)$$

where PX is the tangential component of ϕX and FX is the normal component of ϕX respectively. Similarly, for any vector field N , normal to M , we put

$$\phi N = tN + fN \quad (1.6)$$

where tN and fN are tangential and normal components of ϕN , respectively.

2. Doubly Warped Products

Doubly warped products can be considered as a generalization of warped product (M, g) is a product manifold of the form $M =_f B \times_b F$ with the metric $g = f^2 g_B + b^2 g_F$, where $b : B \rightarrow (0, \infty)$ and $f : F \rightarrow (0, \infty)$ are smooth maps and g_B, g_F are the metrics on the Riemannian manifolds B and F respectively [9]. If either $b = 1$ or $f = 1$, but not both, then we obtain a (*single*) warped product. If both $b = 1$ and $f = 1$, then we have a *product manifold*. If neither b nor f is constant, then we have a non *trivial doubly warped product*.

If $X \in T(B)$ and $Z \in T(F)$, then the Levi-Civita connection is

$$\nabla_X Z = Z(\ln f)X + X(\ln b)Z \tag{2.1}$$

Let $M = {}_{f_2}N_{\perp} \times {}_{f_1}N_T$ be doubly warped product CR-submanifolds of Kenmotsu manifold \bar{M} . Such submanifolds are always tangent to the structure vector field ξ .

Theorem 2.1. *There is no proper doubly warped product contact CR-submanifolds of Kenmotsu manifold \bar{M} .*

Proof. Let $M = {}_{f_2}N_{\perp} \times {}_{f_1}N_T$ be doubly warped product contact CR-submanifolds of Kenmotsu manifold \bar{M} . There are two possible case arise here.

Case (i): When ξ tangent to N_T , then let $Z \in TN_{\perp}$

$$\nabla_Z \xi = Z(\ln f_1)\xi + \xi(\ln f_2)Z \tag{2.2}$$

Also, by equations (1.1) and (1.2)

$$\bar{\nabla}_Z \xi = Z - \eta(Z)\xi = Z$$

or

$$\nabla_Z \xi + h(Z, \xi) = Z$$

this means that

$$\left. \begin{aligned} \nabla_Z \xi &= Z, \\ h(Z, \xi) &= 0. \end{aligned} \right\} \tag{2.3}$$

Using equations (2.2) and (2.3), we get

$$Z = Z(\ln f_1)\xi + \xi(\ln f_2)Z.$$

By the orthogonality of two distribution, we get

$$\left. \begin{aligned} Z(\ln f_1) &= 0, \\ \xi(\ln f_2) &= 1. \end{aligned} \right\} \tag{2.4}$$

First part of equation (2.4) shows that f_2 is constant for all $Z \in TN_{\perp}$. In this case doubly warped product does not exists.

Case (ii) : When ξ tangent to N_{\perp} and let $X \in TN_T$

$$\nabla_X \xi = \xi(\ln f_1)X + X(\ln f_2)\xi \quad (2.5)$$

From structure equation and Gauss formula, we get

$$X - \eta(X)\xi = \bar{\nabla}_X \xi = \nabla_X \xi + h(X, \xi)$$

i.e.,

$$\left. \begin{aligned} \nabla_X \xi &= X, \\ h(X, \xi) &= 0. \end{aligned} \right\} \quad (2.6)$$

Equations (2.5) and (2.6) gives

$$X = (\xi \ln f_1)X + (X \ln f_2)\xi \quad (2.7).$$

By orthogonality of two distributions, equation (2.7) gives

$$\left. \begin{aligned} \xi(\ln f_1) &= 1, \\ X(\ln f_2) &= 0. \end{aligned} \right\} \quad (2.8)$$

Second part of equation (2.8) follows that f_2 is constant for all $X \in T(N_T)$. In this case also, doubly warped product does not exist. Hence the proof is complete.

As in both the cases doubly warped product submanifold $M = {}_{f_2}N_{\perp} \times {}_{f_1}N_T$, ξ tangent to N_T and ξ tangent to N_{\perp} doubly warped products does not exist. Now, we study warped product submanifolds.

3. Warped Product Submanifolds

B.Y. Chen studied warped product CR-submanifolds in Kaehler manifolds and introduce the notion of CR-warped product [4]. Later I. Hasegawa and I. Mihai studied contact CR-warped product submanifolds in Sasakian manifolds [5]. In this section we study contact warped product CR-submanifolds of Kenmotsu manifolds.

Definition 3.1. Let (B, g_1) and (F, g_2) be two Riemannian manifolds with Riemannian metric g_1 and g_2 respectively and f a positive differentiable

function on B . The warped product of B and F is the Riemannian manifold $B \times_f F = (B \times F, g)$, where

$$g = g_1 + f^2 g_2. \quad (3.1)$$

More explicitly, if U is tangent to $M = B \times_f F$ at (p, q) , then

$$\|U\|^2 = \|d\pi_1 U\|^2 + f^2(p) \|d\pi_2 U\|^2$$

where $\pi_i (i = 1, 2)$ are the canonical projections of $B \times F$ onto B and F respectively.

We recall that on a warped product one has

$$\nabla_U V = \nabla_V U = (U \ln f) V \quad (3.2)$$

for any vector fields U tangent to B and V tangent to F [3].

Let $M = N_\perp \times_f N_T$ be a contact warped product CR-submanifolds of Kenmotsu manifold \bar{M} . Such submanifolds are always tangent to the structure vector field ξ . We distinguish 2 cases

- (i) ξ tangent to N_T .
- (ii) ξ tangent to N_\perp .

First we consider the case (i), when ξ tangent to N_T .

Theorem 3.1. *Let \bar{M} be a $(2m + 1)$ -dimensional Kenmotsu manifold. Then there do not exist warped product submanifolds $M = N_\perp \times_f N_T$ such that N_T is an invariant submanifold tangent to ξ and N_\perp is anti-invariant submanifold of \bar{M} .*

Proof. Assume $M = N_\perp \times_f N_T$ is a contact warped product CR-submanifold of a Kenmotsu manifold \bar{M} , such that N_T is an invariant submanifold tangent to ξ and N_\perp is an anti-invariant submanifold of \bar{M} .

By equation (3.2), we get

$$\nabla_X Z = \nabla_Z X = (Z \ln f) X \quad (3.3)$$

for any vector fields Z and X tangent to N_\perp and N_T respectively.

In particular,

$$\nabla_Z \xi = (Z \ln f) \xi \quad (3.4)$$

Also, by structure equation (1.1) and Gauss formula, we have

$$Z = \bar{\nabla}_Z \xi = \nabla_Z \xi + h(Z, \xi)$$

$$\left. \begin{aligned} \nabla_Z \xi &= Z, \\ h(Z, \xi) &= 0. \end{aligned} \right\} \quad (3.5)$$

Equations (3.4) and (3.5) follows that $Z \ln f = 0, \forall Z \in T(N_\perp)$ i.e., f is constant for all $Z \in T(N_\perp)$. This means that N_\perp is trivial, i.e., M is invariant submanifold. This complete the proof of the theorem.

Now, we consider the case (ii), when ξ tangent to N_\perp .

Assume that \bar{M} be $(2m+1)$ -dimensional Kenmotsu manifold and consider the warped product submanifold $M = N_\perp \times_f N_T$ such that N_\perp is an anti-invariant submanifold of dimension p tangent to ξ and N_T is an invariant submanifold of \bar{M} . Then for any $X \in T(N_T)$

$$g(\nabla_X \xi, X) = g(\bar{\nabla}_X \xi, X) = g(X - \eta(X)\xi, X)$$

i.e.,

$$\xi \ln f \|X\|^2 = \|X\|^2$$

or,

$$\xi \ln f = 1$$

i.e.,

$$g(\nabla \ln f, \xi) = 1$$

i.e.,

$$\nabla \ln f = \xi, \quad (3.9)$$

where $\nabla \ln f$ denotes the gradient of $\ln f$ and defined as $g(\nabla \ln f, U) = U \ln f$, for all $U \in T(M)$. Equation (3.9) can also be written as

$$\sum_{i=1}^p \frac{\partial \ln f_i}{\partial x_i} = \xi, \quad i = 1, 2 \cdots p. \quad (3.10)$$

Equation (3.10) is the first order partial differential equation and has a unique solution, i.e., warped product exist in this case.

Now, we will give an example of warped product contact CR-submanifold of type $N_{\perp} \times_f N_T$ in Kenmotsu manifold, with ξ tangent to N_{\perp} :

Example. Consider the complex space C^8 with the usual Kaehler structure and real global coordinates $(x^1, y^1, \dots, x^8, y^8)$. Let $\bar{M} = R \times_f C^8$ be the warped product between the real line R and C^8 , where the warping function is $f = e^t$, t being the global coordinate on R . Then \bar{M} is Kenmotsu manifold [1]. Consider the distribution $D = \text{span}\{\frac{\partial}{\partial x^1}, \frac{\partial}{\partial y^1}, \dots, \frac{\partial}{\partial x^4}, \frac{\partial}{\partial y^4}\}$ and $D^{\perp} = \text{span}\{\frac{\partial}{\partial t}, \frac{\partial}{\partial x^5}, \dots, \frac{\partial}{\partial x^8}, \}$ which are integrable and denote by N_T and N_{\perp} , the integral submanifolds, respectively. Let $g_{N_T} = \sum_{i=1}^4 ((dx^i)^2 + (dy^i)^2)$ and $g_{N_{\perp}} = dt^2 + e^{2t} \sum_{a=5}^8 (dx^a)^2$ be Riemannian metrics on N_T and N_{\perp} , respectively. Then, $M = N_{\perp} \times_f N_T$ is a contact CR-submanifold, isometrically immersed in \bar{M} . Here, the warping function is $f = e^t$.

References

- [1] D.E. Blair, Riemannian Geometry of Contact and Symplectic manifolds, Progress in Mathematics, Vol. 203, Birkhauser, Boston, Basel, Berlin, (2001).
- [2] R.L. Bishop and B. O'Neill, Manifolds of Negative curvature, *Trans. Amer. Math. Soc.*, **145**, (1969) 1-49.
- [3] B.Y. Chen, On isometric minimal immersions from warped products into real space forms, *Proc. Edinburg Math. Soc.* **45** (2002), 579-587.
- [4] B.Y. Chen, Geometry of warped product CR-submanifolds in Kaehler manifold, *Monatsh. math.* **133** (2001), 177-195.
- [5] I. Hasegawa, and I. Mihai, Contact CR-warped product submanifolds in Sasakian manifolds, *Geom. Dedicata* **102** (2003), 143-150.
- [6] M.I. Munteanu, A note on doubly warped product contact CR-submanifolds in trans-sasakian manifolds, To appear in Acta Mathematica Hungaria (2007) (eprint: arXiv:math.DG/0604008 v1).

- [7] C. Murathan, K. Arslan, R. Ezentas, I. Mihai, Contact CR-Warped product submanifolds in Kenmotsu space forms, *J. Korean Math. Soc.* **42** (2005), 1101-1110.
- [8] B. Sahin, Nonexistence of warped product semi-slant submanifolds of Kaehler manifold, *Geom. Dedicata* **117** (2006) 195-202.
- [9] B. Unal, Doubly warped products, *Differential geometry and Its Applications* **15** (3) (2001), 253-263.

(Received 30 August 2007)

Viqar Azam Khan, Khalid Ali Khan, Sirajuddin
Department of Mathematics, Aligarh Muslim University
Aligarh-202 002, India
email: viqarster@gmail.com, khalid.mathematics@gmail.com, siraj@math.com