



A Common Fixed Point Theorem in Fuzzy Metric Space using the Property (CLR_g)

Manish Jain^{†,1} and Sanjay Kumar[‡]

[†]Department of Mathematics, Ahir College
Rewari 123401, India

e-mail : manish_261283@rediffmail.com

[‡]Department of Mathematics, DCRUST
Murthal, Sonapat, India

e-mail : sanjuciet@rediffmail.com

Abstract : In this paper, we generalize the results of Kumar and Fisher [S. Kumar, B. Fisher, A common fixed point theorem in fuzzy metric space using property (E.A.) and implicit relation, Thai J. Math. 8 (3) (2010) 439–446.] using weakly compatible mappings along with property (CLR_g). We also provide an example in support our result.

Keywords : fuzzy metric space; common fixed point; weakly compatible maps; implicit relation; property (CLR_g).

2010 Mathematics Subject Classification : 54H25; 47H10.

1 Introduction

It proved a turning point in the development of mathematics when the notion of fuzzy set was introduced by Zadeh [1]. This notion laid the foundation of fuzzy mathematics. Kramosil and Michalek [2] introduced the notion of a fuzzy metric space by generalizing the concept of the probabilistic metric space to the fuzzy situation. George and Veeramani [3] modified the concept of fuzzy metric spaces introduced by Kramosil and Michalek [2]. There are many view points of the notion of the metric space in fuzzy topology for instance one can refer to Kaleva

¹Corresponding author.

and Seikkala [4], Kramosil and Michalek [2] and George and Veeramani [3]. This proved a milestone in fixed point theory of fuzzy metric space and afterwards a flood of papers appeared for fixed point theorems in fuzzy metric space.

Mishra et al. [5] introduced the concept of compatible maps in FM-spaces which was further generalised by Singh and Jain [6] by introducing the notion of weak compatibility in FM-spaces. In 2002, Aamri and Moutawakil [7] introduced property (E.A.), which is a true generalization of non-compatible maps in metric spaces. Common fixed points for a pair of maps under the notion of property (E.A.) and non-compatible maps were studied by Pant and Pant [8]. Recently, Sintunavarat and Kuman [9] introduced a new concept of property (CLRg). The importance of property (CLRg) ensures that one does not require the closeness of range subspaces and hence, now a days, authors are giving much attention to this property for generalizing the results present in the literature. Works noted in the references [10–14] are some examples.

Popa [15, 16] introduced the idea of implicit function to prove a common fixed point theorem in metric spaces. Jain [17] further extended the result of Popa [15, 16] in fuzzy metric spaces. Afterwards, implicit relations are used as a tool for finding common fixed point of contraction maps (see, [18–23]). Altun and Turkoglu [24] proved two common fixed point theorems on complete FM-space with an implicit relation. In [24], common fixed point theorems have been proved for continuous compatible maps of type (α) or (β) . Kumar and Fisher [25] generalized the results of Altun and Turkoglu [24] by removing the assumption of continuity, relaxing compatibility to weak compatibility and replacing the completeness of the space with a set of four alternative conditions for functions satisfying an implicit relation in FM-space. Our aim is to further generalize the result of Kumar and Fisher [25] by using the property (CLRg) and relaxing many conditions involved.

2 Preliminaries

Before we give our main result we need the following definitions:

Definition 2.1 ([1]). A fuzzy set A in X is a function with domain X and values in $[0, 1]$.

Definition 2.2 ([26]). A binary operation $*$: $[0, 1] \times [0, 1] \rightarrow [0, 1]$ is a continuous t -norm if $([0, 1], *)$ is a topological abelian monoid with unit 1 s.t. $a * b \leq c * d$ whenever $a \leq c$ and $b \leq d$ for all $a, b, c, d \in [0, 1]$.

Definition 2.3 ([3]). The 3-tuple $(X, M, *)$ is called a *fuzzy metric space* if X is an arbitrary set, $*$ is a continuous t -norm and M is a fuzzy set on $X^2 \times [0, \infty)$ satisfying the following conditions:

$$(FM-1) \quad M(x, y, 0) > 0,$$

$$(FM-2) \quad M(x, y, t) = 1 \text{ iff } x = y,$$

$$(FM-3) \quad M(x, y, t) = M(y, x, t),$$

(FM-4) $M(x, y, t) * M(y, z, s) \leq M(x, z, t + s),$

(FM-5) $M(x, y, \cdot) : (0, \infty) \rightarrow [0, 1]$ is continuous, for all $x, y, z \in X$ and $s, t > 0$.

Throughout this paper, we consider M to be a fuzzy metric space with condition:

(FM-6) $\lim_{t \rightarrow \infty} M(x, y, t) = 1$ for all $x, y \in X$ and $t > 0$.

Definition 2.4 ([3]). Let $(X, M, *)$ be fuzzy metric space. A sequence $\{x_n\}$ in X is said to be

(i) *Convergent to a point* $x \in X$, if $\lim_{n \rightarrow \infty} M(x_n, x, t) = 1$ for all $t > 0$;

(ii) *Cauchy sequence* if $\lim_{n \rightarrow \infty} M(x_{n+p}, x_n, t) = 1$, for all $t > 0$ and $p > 0$.

Definition 2.5 ([3]). A fuzzy metric space $(X, M, *)$ is said to be *complete* if and only if every Cauchy sequence in X is convergent.

Lemma 2.6 ([27]). $M(x, y, \cdot)$ is non-decreasing for all $x, y \in X$.

Lemma 2.7 ([27]). Let $x_n \rightarrow x$ and $y_n \rightarrow y$, then

(i) $\lim_{n \rightarrow \infty} M(x_n, y_n, t) \geq M(x, y, t)$, for all $t > 0$,

(ii) $\lim_{n \rightarrow \infty} M(x_n, y_n, t) = M(x, y, t)$, for all $t > 0$, if $M(x, y, t)$ is continuous.

Lemma 2.8 ([5]). If for all $x, y \in X$, $t > 0$ and for a number $k \in (0, 1)$;

$$M(x, y, kt) \geq M(x, y, t), \quad \text{then } x = y.$$

Definition 2.9 ([5]). Let A and B be maps from a FM-space (X, M, \cdot) into itself. The maps A and B are said to be *compatible* (or *asymptotically commuting*), if for all t ,

$$\lim_{n \rightarrow \infty} M(AB_{x_n}, BA_{x_n}, t) = 1,$$

whenever $\{x_n\}$ is a sequence in X such that

$$\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Bx_n = z \quad \text{for some } z \in X.$$

From the above definition it is inferred that A and B are non-compatible maps from a FM-space (X, M, \cdot) into itself if $\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Bx_n = z$ for some $z \in X$, but either $\lim_{n \rightarrow \infty} M(AB_{x_n}, BA_{x_n}, t) \neq 1$ or the limit does not exist.

Definition 2.10 ([6]). Let A and B be maps from a FM-space (X, M, \cdot) into itself. The maps are said to be *weakly compatible* if they commute at their coincidence points. Note that compatible mappings are weakly compatible but converse is not true in general.

Definition 2.11 ([8]). Let A and B be two self-maps of a FM-space (X, M, \cdot) . We say that A and B satisfy the property (E.A.) if there exists a sequence $\{x_n\}$ such that

$$\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} Bx_n = z \quad \text{for some } z \in X.$$

Note that weakly compatible and property (E.A.) are independent to each other (see [15], Example 2.2).

Definition 2.12 ([9]). Let (X, d) be a metric space. Two mappings $f : X \rightarrow X$ and $g : X \rightarrow X$ are said to satisfy property (CLR_g) if there exists sequences $\{x_n\}$ in X such that

$$\lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} g(x_n) = g(p), \quad \text{for some } p \text{ in } X.$$

Similarly, we can have the property (CLR_T) and the property (CLR_S) if in the Definition 2.12, the mapping $g : X \rightarrow X$ has been replaced by the mapping $T : X \rightarrow X$ and $S : X \rightarrow X$ respectively.

Our result deal with the following implicit relation used by Altun and Turkoglu [24].

Definition 2.13 ([24]). Let $I = [0, 1]$, $*$ be a continuous t -norm and \mathcal{F} be the set of all real continuous functions $F : I^6 \rightarrow R$ satisfying the following conditions:

(F-1) F is non-increasing in the fifth and sixth variables,

(F-2) if for some constant $k \in (0, 1)$ we have

$$(F-a) \quad F\left(u(kt) \cdot v(t), v(t), u(t), 1, u\left(\frac{t}{2}\right) * v\left(\frac{t}{2}\right)\right) \geq 1,$$

or

$$(F-b) \quad F\left(u(kt), v(t), u(t), v(t), u\left(\frac{t}{2}\right) * v\left(\frac{t}{2}\right), 1\right) \geq 1,$$

for any fixed $t > 0$ and any non-decreasing functions $u, v : (0, \infty) \rightarrow I$, then there exists $h \in (0, 1)$ with $u(ht) \geq v(t) * u(t)$,

(F-3) if for some constant $k \in (0, 1)$, we have $F(u(kt), u(t), 1, 1, u(t), u(t)) \geq 1$ for any fixed $t > 0$ and any non-decreasing function $u : (0, \infty) \rightarrow I$ then $u(kt) \geq u(t)$.

3 Main results

In [24], Altun and Turkoglu proved the following result:

Theorem 3.1. *Let $(X, M, *)$ be a complete fuzzy metric space with $a * b = \min\{a, b\}$. Let A, B, S, T be maps from X into itself satisfying the following conditions:*

- (3.1) $A(X) \subseteq T(X), B(X) \subseteq S(X)$;
- (3.2) *one of the maps A, B, S, T is continuous;*
- (3.3) *the pairs (A, S) and (B, T) are compatible of type (α) ;*
- (3.4) *there exists $k \in (0, 1)$ and $F \in \mathcal{F}$ such that*

$$F\{M(Ax, By, kt), M(Sx, Ty, t), M(Ax, Sx, t), M(By, Ty, t), M(Ax, Ty, t), M(By, Sx, t)\} \geq 1$$

for all $x, y \in X$ and $t > 0$.

Then A, B, S, T have a unique common fixed point in X .

In [25], Kumar and Fisher generalized Theorem 3.1 (which is Theorem 1 in [24]) as follows:

Theorem 3.2. *Let $(X, M, *)$ be a fuzzy metric space with $a * b = \min\{a, b\}$. Further, let (A, S) and (B, T) be weakly compatible pairs of self-maps of X satisfying (3.1), (3.4) with the following condition:*

- (3.5) *one of the pairs (A, S) or (B, T) satisfies property (E.A.).*

If the range of one of the maps A, B, S or T is a complete subspace of X , then A, B, S, T have a unique common fixed point in X .

We now generalize Theorem 3.2 as follows:

Theorem 3.3. *Let $(X, M, *)$ be a fuzzy metric space with $a * b = \min\{a, b\}$. Let A, B, S, T be maps from X into itself satisfying (3.4) with the following conditions:*

- (3.6) *$B(X) \subseteq S(X)$ and the pair (B, T) satisfies property (CLR_T) ,*
or
 $A(X) \subseteq T(X)$ and the pair (A, S) satisfies property (CLR_S) ;
- (3.7) *the pairs (A, S) and (B, T) are weakly compatible.*

Then A, B, S, T have a unique common fixed point in X .

Proof. Without loss of generality, assume that $B(X) \subseteq S(X)$ and the pair (B, T) satisfies property (CLR_T) , then there exists a sequence $\{x_n\}$ in X such that Bx_n and Tx_n converges to Tx , for some x in X as $n \rightarrow \infty$. Since $B(X) \subseteq S(X)$, so there exists a sequence $\{y_n\}$ in X such that $Bx_n = Sy_n$, hence $Sy_n \rightarrow Tx$ as $n \rightarrow \infty$.

We shall show that $\lim_{n \rightarrow \infty} Ay_n = Tx$. Let $\lim_{n \rightarrow \infty} Ay_n = z$. Taking $x = y_n, y = x_n$ in (3.4),

$$F\{M(Ay_n, Bx_n, kt), M(Sy_n, Tx_n, t), M(Ay_n, Sy_n, t), M(Bx_n, Tx_n, t), M(Ay_n, Tx_n, t), M(Bx_n, Sy_n, t)\} \geq 1.$$

Letting $n \rightarrow \infty$, we have

$$F\{M(z, Tx, kt), 1, M(z, Tx, t), 1, M(z, Tx, t), 1\} \geq 1.$$

On the other hand, since

$$M(z, Tx, t) \geq M\left(z, Tx, \frac{t}{2}\right) = M\left(z, Tx, \frac{t}{2}\right) * 1,$$

and F is non-increasing in the fifth variable, we have, for any $t > 0$

$$\begin{aligned} F\left\{M(z, Tx, kt), 1, M(z, Tx, t), 1, M\left(z, Tx, \frac{t}{2}\right), 1\right\} \\ \geq F\{M(z, Tx, kt), 1, M(z, Tx, t), 1, M(z, Tx, t)\} \geq 1, \end{aligned}$$

which implies by (F-2), that $z = Tx$. Subsequently, we have Bx_n, Tx_n, Sy_n, Ay_n converges to z . We shall show that $Bx = z$.

Taking $x = y_n, y = x$ in (3.4),

$$\begin{aligned} F\{M(Ay_n, Bx, kt), M(Sy_n, Tx, t), M(Ay_n, Sy_n, t), M(Bx, Tx, t), \\ M(Ay_n, Tx, t), M(Bx, Sy_n, t)\} \geq 1. \end{aligned}$$

Letting $n \rightarrow \infty$, we have

$$F\{M(z, Bx, kt), 1, 1, M(z, Bx, t), 1, M(z, Bx, t)\} \geq 1.$$

On the other hand, since

$$M(z, Bx, t) \geq N\left(z, Bx, \frac{t}{2}\right) = M\left(z, Bx, \frac{t}{2}\right) * 1,$$

and F is non-increasing in the sixth variable, we have, for any $t > 0$

$$\begin{aligned} F\left\{M(z, Bx, kt), 1, 1, M(z, Bx, t), 1, M\left(z, Bx, \frac{t}{2}\right) * 1\right\} \\ \geq F\{M(z, Bx, kt), 1, 1, M(z, Bx, t), 1, M(z, Bx, t)\} \geq 1, \end{aligned}$$

which implies by (F-2) that $z = Bx = Tx$. Since, the pair (B, T) is weak compatible, it follows that $Bz = Tz$.

Also, since $B(X) \subseteq S(X)$, there exists some y in X such that $Bx = Sy (= z)$.

We next show that $Sy = Ay (= z)$. Taking $y = x_n, x = y$ in (3.4),

$$\begin{aligned} F\{M(Ay, Bx_n, kt), M(Sy, Tx_n, t), M(Ay, Sy, t), M(Bx_n, Tx_n, t), \\ M(Ay, Tx_n, t), M(Bx_n, Sy, t)\} \geq 1. \end{aligned}$$

Letting $n \rightarrow \infty$, we have

$$F\{M(Ay, z, kt), 1, M(Ay, z, t), 1, M(Ay, z, t), 1\} \geq 1.$$

Other the other hand, since

$$M(Ay, z, t) \geq M\left(Ay, z, \frac{t}{2}\right) = M\left(Ay, z, \frac{t}{2}\right) * 1,$$

and F is non-increasing in the fifth variable, we have, for any $t > 0$

$$F\left\{M(Ay, z, kt), 1, M(Ay, z, t), 1, M\left(Ay, z, \frac{t}{2}\right) * 1, 1\right\} \\ \geq F\{M(Ay, z, kt), 1, M(Ay, z, t), 1, M(Ay, z, t), 1\} \geq 1,$$

which implies by (F-2) that $Ay = z = Sy$. But the pair (A, S) is weakly compatible, it follows that $Az = Sz$.

Next, we claim that $Az = Bz$. Taking $x = z, y = z$ in (3.4),

$$F\{M(Az, Bz, kt), M(Az, Bz, t), 1, 1, M(Az, Bz, t), M(Az, Bz, t)\} \geq 1,$$

which implies by (F-3) that $Az = Bz$. Hence, $Az = Bz = Sz = Tz$.

We now show that $z = Az$. Taking $x = z, y = x$ in (3.4),

$$F\{M(Az, Bx, kt), M(Sz, Tx, t), M(Az, Sz, t), M(Bx, Tx, t), \\ M(Az, Tx, t), M(Bx, Sz, t)\} \geq 1,$$

that is,

$$F\{M(Az, z, kt), M(Az, z, t), 1, 1, M(Az, z, t), M(Az, z, t)\} \geq 1.$$

Therefore, $z = Az = Bz = Sz = Tz$, that is z is the common fixed point of the maps A, B, S, T . Uniqueness of z follows immediately from (F-3) and (3.4). \square

Example 3.4. Let $(X, M, *)$ be a fuzzy metric space with $X = [0, 1]$, a t -norm $*$ be defined by $a * b = \min\{a, b\}$ for all a, b in $[0, 1]$ and M be a fuzzy set on $X^2 \times (0, \infty)$ defined by

$$M(x, y, t) = \left[\exp\left(\frac{|x - y|}{t}\right) \right]^{-1}$$

for all x, y in X and $t > 0$.

Let $F : I^6 \rightarrow R$ be defiend by $F(u_1, u_2, u_3, u_4, u_5, u_6) = \frac{u_1}{\min\{u_2, u_3, u_4, u_5, u_6\}}$. Let $t > 0, 0 < u(t), v(t) \leq 1, k \in (0, \frac{1}{2})$, where $u, v : [0, \infty) \rightarrow I$ are non-decreasing functions. Suppose that

$$F\left(u(kt), v(t), v(t)u(t), 1, u\left(\frac{t}{2}\right) * v\left(\frac{t}{2}\right)\right) \geq 1,$$

that is,

$$F\left(u(kt), v(t), v(t), u(t), 1, u\left(\frac{t}{2}\right) * v\left(\frac{t}{2}\right)\right) = \frac{u(kt)}{\min\{v(t), v(t), u(t), 1, u\left(\frac{t}{2}\right) * v\left(\frac{t}{2}\right)\}} \\ \geq 1.$$

Thus $u(ht) \geq v(t) * u(t)$ if $h = 2k \in (0, 1)$. A similar argument works if (F_b) is assumed. Finally, suppose that $t > 0$ is fixed, $u : [0, \infty) \rightarrow I$ is a non-decreasing function and

$$F(u(kt), u(t), 1, 1u(t), u(t)) = \frac{u(kt)}{u(t)} \geq 1,$$

for some $k \in (0, 1)$. Then we have $u(kt) \geq u(t)$ and thus $F \in \mathcal{F}$.

Define the mappings $A, B, S, T : X \rightarrow X$ by

$$Ax = \frac{x}{27}, \quad Bx = \frac{x}{9}, \quad Sx = \frac{x}{3}, \quad Tx = x,$$

respectively. Then, for some $k \in [\frac{1}{9}, 1)$, we have

$$\begin{aligned} M(Ax, By, kt) &= \left[\exp \left(\frac{|\frac{x}{27} - \frac{y}{9}|}{kt} \right) \right]^{-1} \\ &\geq \left[\exp \left(\frac{|\frac{x}{3} - y|}{t} \right) \right]^{-1} \\ &= M(Sx, Ty, t) \\ &\geq \min\{M(Sx, Ty, t), M(Ax, Sx, t), M(By, Ty, t), M(Ax, Ty, t), \\ &\quad M(By, Sx, t)\}. \end{aligned}$$

Thus, the condition (3.4) of Theorem 3.3 is satisfied.

Further, the pairs (A, S) and (B, T) are weakly compatible. Also, $B(X) = [0, \frac{1}{9}] \subseteq [0, \frac{1}{3}] = S(X)$. Considering the sequence $\{x_n\} = \{\frac{1}{n}\}$ so that $\lim_{n \rightarrow \infty} Bx_n = \lim_{n \rightarrow \infty} Tx_n = 0 = T(0)$, hence the pair (B, T) satisfies property (CLR_T) .

Therefore, all the conditions of Theorem 3.3 are satisfied. Indeed 0 is the unique common fixed point of the mappings A, B, S, T .

References

- [1] L.A. Zadeh, Fuzzy sets, Infor. Control 8 (1965) 338–353.
- [2] I. Kramosil, J. Michalek, Fuzzy metric and statistical metric spaces, Kybernetika 11 (1975) 336–344.
- [3] A. George, P. Veeramani, On some results in fuzzy metric spaces, Fuzzy Sets and Systems 64 (1994) 395–399.
- [4] O. Kaleva, S. Seikkala. On fuzzy metric spaces, Fuzzy Sets and Systems 12 (3) (1984) 215–229.
- [5] S.N. Mishra, N. Sharma, S.L. Singh, Common fixed point of maps on fuzzy metric spaces, Internat. J. Math. Math. Sci. 17 (1994) 253–258.
- [6] B. Singh, S. Jain, Weak compatibility and fixed point theorems in fuzzy metric spaces, Ganita 56 (2) (2005) 167–176.

- [7] M. Aamri, D. El Moutawakil, Some new common fixed point theorems under strict contractive conditions, *J. Math. Anal. Appl.* 270 (2002) 181–188.
- [8] V. Pant, R.P. Pant, Fixed points in fuzzy metric space for noncompatible maps, *Soochow J. Math.* 33 (4) (2007) 647–655.
- [9] W. Sintunavarat, P. Kumam, Common fixed point theorems for a pair of weakly compatible mappings in Fuzzy Metric Spaces, *Journal of Applied Mathematics*, Vol. 2011 (2011), Article ID 637958, 14 pages.
- [10] W. Sintunavarat, P. Kumam, Common fixed points for R-weakly commuting in fuzzy metric spaces, *Annali dell' Universit'a di Ferrara*, (In Press).
- [11] S. Chauhan, W. Sintunavarat, P. Kumam, Common fixed point theorems for weakly compatible mappings in fuzzy metric spaces using (JCLR) property, *Applied Mathematics*, (to appear).
- [12] M. Imdad, B.D. Pant, S. Chauhan, Fixed point theorems in menger spaces using the (CLR_{ST}) property and applications, *Journal of Nonlinear Analysis and Optimization: Theory & Applications*, (In Press).
- [13] S.L. Singh, B.D. Pant, S. Chauhan, Fixed point theorems in non-archimedean menger PM-spaces, *Journal of Nonlinear Analysis and Optimization: Theory & Applications*, (In Press).
- [14] M. Jain, K. Tas, S. Kumar, N. Gupta, Coupled fixed point theorems for a pair of weakly compatible maps along with CLRg property in fuzzy metric spaces, *Journal of Applied Mathematics*, (In Press).
- [15] V. Popa, Some fixed point theorems for compatible mappings satisfying on implicit relation, *Demonstratio Mathematica* 32 (1999) 157–163.
- [16] V. Popa, A general coincidence theorem for compatible multivalued mappings satisfying an implicit relation, *Demonstratio Mathematica* 33 (2000) 159–164.
- [17] S. Jain, B. Mundra, S. Aake, Common fixed point theorem in fuzzy metric space using implicit relation, *Int. Math. Forum* 4 (3) (2009) 135–141.
- [18] A. Aliouche, V. Popa, Common fixed point theorems for occasionally weakly compatible mappings via implicit relations, *Filomat* 22 (2) (2008) 99–107.
- [19] S. Kumar, B.D. Pant, A common fixed point theorem in probabilistic metric space using implicit relation, *Filomat* 22 (2) (2008) 43–52.
- [20] H.K. Pathak, R.R. Lopez, R.K. Verma, A common fixed point theorem using implicit relation and property (E.A) in metric spaces, *Filomat* 21 (2) (2007) 211–234.
- [21] V. Popa, A generalization of Meir-Keeler type common fixed point theorem for four noncontinuous mappings, *Sarajevo J. Math.* 1 (13) (2005) 135–142.

- [22] S. Sedghi, K.P.R. Rao, N. Shobe, A general common fixed point theorem for multi-maps satisfying an implicit relation on fuzzy metric spaces, *Filomat* 22 (1) (2008) 1–10.
- [23] B. Singh, S. Jain, Semicompatibility and fixed point theorems in fuzzy metric space using implicit relation, *Int. J. Math. Math. Sci.* 2005 (16) (2005) 2617–2629.
- [24] I. Altun, D. Turkoglu, Some fixed point theorems on fuzzy metric spaces with implicit relations, *Commun. Korean Math. Soc.* 23 (1) (2008) 111–124.
- [25] S. Kumar, B. Fisher, A common fixed point theorem in fuzzy metric space using property (E.A.) and implicit relation, *Thai J. Math.* 8 (3) (2010) 439–446.
- [26] B. Schweizer, A. Sklar, *Probabilistic Metric Spaces*, North Holland Series in Probability and Applied Math., Amsterdam, vol. 5, (1983).
- [27] M. Grabiec, Fixed points in fuzzy metric spaces, *Fuzzy Sets and Systems* 27 (3) (1988) 385–389.

(Received 31 July 2012)

(Accepted 27 August 2012)