



Projection Type Ishikawa Iteration with Perturbations for Common Fixed Points of Two Nonself Generalized Asymptotically Quasi-Nonexpansive Mappings

Kritsadaphiwat Wongyai and Tanakit Thianwan¹

School of Science, University of Phayao, Phayao 56000, Thailand

e-mail : kritsadaphiwat@gmail.com (K. Wongyai)

tanakit.th@up.ac.th (T. Thianwan)

Abstract : In this paper, we introduce and study a new type of two-step iterative scheme which is called the projection type Ishikawa iteration with perturbations for two nonself generalized asymptotically quasi-nonexpansive mappings in Banach spaces. A sufficient condition for convergence of the iteration process to a common fixed point of mappings under our setting is also established in a real uniformly convex Banach space. Furthermore, the strong convergence of a new iterative scheme with perturbations to a common fixed point of two nonself generalized asymptotically quasi-nonexpansive mappings on a nonempty closed convex subset of a real Banach space is proved. The results obtained in this paper extend and generalize many important know results in recent literature.

Keywords : nonself generalized asymptotically quasi-nonexpansive mapping; uniformly L -Lipschitzian; strong convergence; completely continuous; common fixed points.

2010 Mathematics Subject Classification : 47H10; 47H09; 46B20.

⁰This research was supported by the revenue budget School of Science, University of Phayao.

¹Corresponding author.

1 Introduction and Preliminaries

As we know, iteration methods are numerical procedures which compute a sequence of gradually accurate iterates to approximate the solution of a class of problems. Such methods are useful tools of applied mathematics for solving real life problems ranging from economics and finance or biology to transportation, network analysis or optimization. When we design iteration methods, we have to study their qualitative properties such as: convergence, stability, error propagation, stopping criteria. This is an active area of research, several well known scientists in the world paid and still pay attention to the qualitative study of iteration methods; please, see [1–4].

Fixed-point iteration process for nonexpansive self-mappings including Mann and Ishikawa iteration processes have been studied extensively by various authors (see [5–7]). We know that Mann and Ishikawa iteration processes are defined as:

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T x_n, \quad n \geq 1, \quad (1.1)$$

and

$$\begin{aligned} y_n &= (1 - \beta_n)x_n + \beta_n T x_n, \\ x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n T y_n, \quad n \geq 1, \end{aligned} \quad (1.2)$$

respectively. Obviously the iterative schemes (1.1) and (1.2) deals with one self-mapping only. In 1986, Das and Debata [6] introduced and studied the case of two mapping in iteration processes. This success can be rich source of inspiration for many authors, see for example, Takahashi and Tamura [8] and Khan and Takahashi [9]. For approximating the common fixed points, the two mappings case has its own importance as it has a direct link with the minimization problem, see for example Takahashi [10].

Being an important generalization of the class of nonexpansive self-mappings, in 1972, Goebel and Kirk [11] introduced the class of asymptotically nonexpansive self-mappings, who proved that if C is a nonempty closed convex subset of a real uniformly convex Banach space and T is an asymptotically nonexpansive self-mapping on C , then T has a fixed point.

In 1991, Schu [12] introduced the following modified Mann iteration process:

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n, \quad n \geq 1, \quad (1.3)$$

to approximate fixed points of asymptotically nonexpansive self-mappings in Hilbert space. Since then, Schu's iteration process has been widely used to approximate fixed points of asymptotically nonexpansive self-mappings in Hilbert space or Banach spaces (see [12–15]).

Let C be a nonempty closed convex subset of real normed linear space X . A self-mapping $T : C \rightarrow C$ is said to be nonexpansive if $\|T(x) - T(y)\| \leq \|x - y\|$ for all $x, y \in C$. A self-mapping $T : C \rightarrow C$ is called asymptotically nonexpansive if there exists a sequence $\{k_n\} \subset [1, \infty)$, $k_n \rightarrow 1$ as $n \rightarrow \infty$ such that

$$\|T^n x - T^n y\| \leq k_n \|x - y\| \quad (1.4)$$

for all $x, y \in C$ and $n \geq 1$. A mapping $T : C \rightarrow C$ is said to be uniformly L -Lipschitzian if there exists a constant $L > 0$ such that

$$\|T^n x - T^n y\| \leq L\|x - y\| \tag{1.5}$$

for all $x, y \in C$ and $n \geq 1$.

It is easy to see that if T is an asymptotically nonexpansive, then it is uniformly L -Lipschitzian with the uniform Lipschitz constant $L = \sup\{k_n : n \geq 1\}$.

Definition 1.1 (see [16]). A self-mapping $T : C \rightarrow C$ is called *generalized asymptotically nonexpansive* if there exists nonnegative real sequences $\{k_n\}$ and $\{\delta_n\}$ with $k_n > 1$, $k_n \rightarrow 1$ and $\delta_n \rightarrow 0$ as $n \rightarrow \infty$ such that

$$\|T^n x - T^n y\| \leq k_n\|x - y\| + \delta_n \tag{1.6}$$

for all $x, y \in C$ and $n \geq 1$. $T : C \rightarrow C$ is said to be *generalized asymptotically quasi-nonexpansive* if there exists nonnegative real sequences $\{k_n\}$ and $\{\delta_n\}$ with $k_n > 1$, $k_n \rightarrow 1$ and $\delta_n \rightarrow 0$ as $n \rightarrow \infty$ such that

$$\|T^n x - T^n p\| \leq k_n\|x - p\| + \delta_n \tag{1.7}$$

for all $x \in C$, $p \in F(T)$ ($F(T)$ denote the set of fixed points of T) and $n \geq 1$.

It is clear from the definition that a generalized asymptotically quasi-nonexpansive mapping is to unify various definitions of classes of mappings associated with the class of generalized asymptotically nonexpansive mapping, asymptotically nonexpansive type, asymptotically nonexpansive mappings, and nonexpansive mappings. However, the converse of each of above statement may be not true. The example shows that a generalized asymptotically quasi-nonexpansive mapping is not an asymptotically quasi-nonexpansive mapping; see [16].

Iterative techniques for approximating fixed points of nonexpansive mappings and their generalizations, for example, asymptotically nonexpansive mappings, etc., have been studied by a number of authors (see, e.g., [13–17]) and references cited therein.

In most of these papers, the well known Mann iteration process (1.1) (see [17]) has been studied and the operator T has been assumed to map C into itself. The convexity of C then ensures that the sequence $\{x_n\}$ generated by (1.1) is well defined. If, however, C is a proper subset of the real Banach space X and T maps C into X (as is the case in many applications), then the sequence given by (1.1) may not be well defined. One method that has been used to overcome this in the case of single operator T is to introduce a retraction $P : X \rightarrow C$ in the recursion formula (1.1) as follows: $x_1 \in C$,

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n P T x_n, \quad n \geq 1. \tag{1.8}$$

For nonself nonexpansive mappings, some authors (see [18–22]) have studied the strong and weak convergence theorems in Hilbert space or uniformly convex Banach space.

The concept of nonself asymptotically nonexpansive mappings was introduced by Chidume, Ofoedu and Zegeye [2] in 2003 as the generalization of asymptotically nonexpansive self-mappings. The nonself asymptotically nonexpansive mapping is defined as follows:

Definition 1.2 (see [2]). Let C be a nonempty subset of a real normed linear space X . Let $P : X \rightarrow C$ be a nonexpansive retraction of X onto C . A nonself-mapping $T : C \rightarrow X$ is called *asymptotically nonexpansive* if there exists a sequence $\{k_n\} \subset [1, \infty)$, $k_n \rightarrow 1$ as $n \rightarrow \infty$ such that

$$\|T(PT)^{n-1}x - T(PT)^{n-1}y\| \leq k_n \|x - y\| \quad (1.9)$$

for all $x, y \in C$ and $n \geq 1$. T is said to be *uniformly L -Lipschitzian* if there exists a constant $L > 0$ such that

$$\|T(PT)^{n-1}x - T(PT)^{n-1}y\| \leq L \|x - y\| \quad (1.10)$$

for all $x, y \in C$ and $n \geq 1$.

In [2], they studied the following iterative sequence: $x_1 \in C$,

$$x_{n+1} = P((1 - \alpha_n)x_n + \alpha_n T(PT)^{n-1}x_n) \quad (1.11)$$

to approximate some fixed point of T under suitable conditions.

If T is a self-mapping, then P becomes the identity mapping so that (1.9) and (1.10) reduce to (1.4) and (1.5), respectively. (1.11) reduces to (1.3).

In 2006, Wang [21] generalizes the iteration process (1.11) as follows: $x_1 \in C$,

$$\begin{aligned} y_n &= P((1 - \beta_n)x_n + \beta_n T_2(PT_2)^{n-1}x_n), \\ x_{n+1} &= P((1 - \alpha_n)x_n + \alpha_n T_1(PT_1)^{n-1}y_n), \quad n \geq 1, \end{aligned} \quad (1.12)$$

where $T_1, T_2 : C \rightarrow X$ are nonself asymptotically nonexpansive mappings and $\{\alpha_n\}, \{\beta_n\}$ are real sequences in $[0, 1)$. He proved strong convergence of the sequence $\{x_n\}$ defined by (1.12) to a common fixed point of T_1 and T_2 under proper conditions. Meanwhile, the results of [21] generalized the results of [2].

The nonself generalized asymptotically nonexpansive and nonself generalized asymptotically quasi-nonexpansive mappings are defined as follows:

Definition 1.3 (see [22]). Let C be a nonempty subset of a real normed linear space X . Let $P : X \rightarrow C$ be a nonexpansive retraction of X onto C . A nonself-mapping $T : C \rightarrow X$ is called *generalized asymptotically nonexpansive* if there exists nonnegative real sequences $\{k_n\}$ and $\{\delta_n\}$ with $k_n > 1$, $k_n \rightarrow 1$ and $\delta_n \rightarrow 0$ as $n \rightarrow \infty$ such that

$$\|T(PT)^{n-1}x - T(PT)^{n-1}y\| \leq k_n \|x - y\| + \delta_n \quad (1.13)$$

for all $x, y \in C$ and $n \geq 1$. $T : C \rightarrow X$ is said to be *generalized asymptotically quasi-nonexpansive* if there exists nonnegative real sequences $\{k_n\}$ and $\{\delta_n\}$ with $k_n > 1$, $k_n \rightarrow 1$ and $\delta_n \rightarrow 0$ as $n \rightarrow \infty$ such that

$$\|T(PT)^{n-1}x - T(PT)^{n-1}p\| \leq k_n\|x - p\| + \delta_n \tag{1.14}$$

for all $x \in C$, $p \in F(T)$ and $n \geq 1$.

If T is a self-mapping, then P becomes the identity mappings so that (1.13) and (1.14) reduces to (1.6) and (1.7), respectively.

In 2008, Deng and Liu [22] studied the following iterative sequence which can be viewed as an extension for iterative schemes of Wang [21]: $x_i \in C$ ($i = 0, 1, 2, \dots, q$ and $q \in \mathbb{N}$ is a fixed number),

$$\begin{aligned} y_n &= P(\bar{\alpha}_n x_n + \bar{\beta}_n T_2(PT_2)^{n-1}x_n + \bar{\gamma}_n v_n), \quad n = 0, 1, 2, \dots, \\ x_{n+1} &= P(\alpha_n x_n + \beta_n T_1(PT_1)^{n-1}y_{n-q} + \gamma_n u_n), \quad n = q, q + 1, q + 2, \dots, \end{aligned} \tag{1.15}$$

where $T_1, T_2 : C \rightarrow X$ are nonself generalized asymptotically quasi-nonexpansive mappings, $\{u_n\}, \{v_n\}$ are bounded sequences in C and $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\bar{\alpha}_n\}, \{\bar{\beta}_n\}$ and $\{\bar{\gamma}_n\}$ are real sequences in $[0, 1]$ satisfying $\alpha_n + \beta_n + \gamma_n = \bar{\alpha}_n + \bar{\beta}_n + \bar{\gamma}_n = 1$ for all $n \geq 0$. They gave the following strong convergence theorem.

Theorem 1.4 (see [22]). *Let X be a real uniformly convex Banach space, C a nonempty closed convex subset of X , $T_1, T_2 : C \rightarrow X$ two uniformly L -Lipschitzian, nonself generalized asymptotically quasi-nonexpansive mappings with nonnegative real sequences $\{k_n^{(i)}\}, \{\delta_n^{(i)}\} \subset [1, \infty)$ ($i = 1, 2$), respectively such that $\sum_{n=1}^{\infty} (k_n^{(i)} - 1) < \infty, \sum_{n=1}^{\infty} \delta_n^{(i)} < \infty$. Suppose $F = F(T_1) \cap F(T_2) \neq \emptyset$. For any $x_i \in C$ ($i = 0, 1, 2, \dots, q$ and $q \in \mathbb{N}$ is a fixed number), let $\{x_n\}$ be the sequence defined by (1.15) satisfying $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1, 0 < \liminf_{n \rightarrow \infty} \bar{\alpha}_n \leq \limsup_{n \rightarrow \infty} \bar{\alpha}_n < 1, \sum_{n=1}^{\infty} \gamma_n < \infty$ and $\sum_{n=1}^{\infty} \bar{\gamma}_n < \infty$. If T_1, T_2 satisfies condition A' with respect to the sequence $\{x_n\}$, then $\{x_n\}$ converges strongly to a common fixed point of T_1, T_2 .*

Recently, a new iterative scheme which is called the projection type Ishikawa iteration for two nonself asymptotically nonexpansive mappings was defined and constructed by Thianwan [23]. It is given as follows:

$$\begin{aligned} y_n &= P((1 - \beta_n)x_n + \beta_n T_2(PT_2)^{n-1}x_n), \\ x_{n+1} &= P((1 - \alpha_n)y_n + \alpha_n T_1(PT_1)^{n-1}y_n), \quad n \geq 1, \end{aligned} \tag{1.16}$$

where $\{\alpha_n\}$ and $\{\beta_n\}$ are appropriate real sequences in $[0, 1]$. He studied the scheme for two nonself asymptotically nonexpansive mappings and proved strong convergence of the sequences $\{x_n\}$ and $\{y_n\}$ to a common fixed point of T_1, T_2 under suitable conditions in a uniformly convex Banach space.

Note that Thianwan process (1.16) and Wang process (1.12) are independent: neither reduces to the other.

If $T_1 = T_2$ and $\beta_n = 0$ for all $n \geq 1$, then (1.16) reduces to (1.11). It also can be reduces to Schu process (1.3).

We note that, in applications, there are perturbations always occurring in the iterative processes because the manipulations are inaccurate. It is no doubt that researching the convergent problems of iterative methods with perturbation members is a significant job. This leads us, in this paper, to introduce and study a new class of two-step iterative scheme with perturbations for solving the fixed point problem for nonself generalized asymptotically quasi-nonexpansive mappings. This iterative scheme can be viewed as an extension for Ishikawa type iterative schemes of Thianwan [23]. The scheme is defined as follows.

Let X be a normed space, C a nonempty convex subset of X , $P : X \rightarrow C$ a nonexpansive retraction of X onto C and $T_1, T_2 : C \rightarrow X$ are given mappings. Then for an arbitrary $x_1 \in C$, the following iteration scheme is studied:

$$\begin{aligned} y_n &= P((1 - \beta_n - \gamma_n)x_n + \beta_n T_2 (PT_2)^{n-1} x_n + \gamma_n v_n), \\ x_{n+1} &= P((1 - \alpha_n - \lambda_n)y_n + \alpha_n T_1 (PT_1)^{n-1} y_n + \lambda_n u_n), \quad n \geq 1, \end{aligned} \quad (1.17)$$

where $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$ and $\{\lambda_n\}$ are appropriate real sequences in $[0, 1]$ and $\{u_n\}, \{v_n\}$ are bounded sequences in C . We then prove its strong convergence under some suitable conditions in Banach spaces.

Note that Deng and Liu process (1.15) and our process (1.17) are independent: neither reduces to the other.

If $\gamma_n = \lambda_n = 0$ for all $n \geq 1$, then (1.17) reduces to (1.16).

Now, we recall some well known concepts and results.

Let X be a Banach space with dimension $X \geq 2$. The modulus of X is the function $\delta_X : (0, 2] \rightarrow [0, 1]$ defined by

$$\delta_X(\epsilon) = \inf \left\{ 1 - \left\| \frac{1}{2}(x + y) \right\| : \|x\| = 1, \|y\| = 1, \epsilon = \|x - y\| \right\}.$$

Banach space X is uniformly convex if and only if $\delta_X(\epsilon) > 0$ for all $\epsilon \in (0, 2]$.

A subset C of X is said to be a retract if there exists a continuous mapping $P : X \rightarrow C$ such that $Px = x$ for all $x \in C$. Every closed convex subset of a uniformly convex Banach space is a retract. A mapping $P : X \rightarrow X$ is said to be a retraction if $P^2 = P$. It follows that if a mapping P is a retraction, then $Pz = z$ for every $z \in R(P)$, the range of P .

A set C is optimal if each point outside C can be moved to be closer to all points of C . It is well known (see [7]) that

(1) If X is a separable, strictly convex, smooth, reflexive Banach space, and if $C \subset X$ is an optimal set with interior, then C is a nonexpansive retract of X .

(2) A subset of l^p , with $1 < p < \infty$, is a nonexpansive retract if and only if it is optimal.

Note that every nonexpansive retract is optimal. In strictly convex Banach spaces, optimal sets are closed and convex. Moreover, every closed convex subset of a Hilbert space is optimal and also a nonexpansive retract.

Recall that two mappings $S, T : C \rightarrow X$, where C is a subset of a normed space X , are said to satisfy condition A' (see [24]) if there exists a nondecreasing

function $f : [0, \infty) \rightarrow [0, \infty)$ with $f(0) = 0$, $f(r) > 0$ for all $r \in (0, \infty)$ such that either

$$\|x - Sx\| \geq f(d(x, F)) \quad \text{or} \quad \|x - Tx\| \geq f(d(x, F))$$

for all $x \in C$, where $d(x, F) = \inf\{\|x - q\| : q \in F = F(S) \cap F(T)\}$.

Note that condition A' reduces to condition (A) (see [15]) when $S = T$. Maiti and Ghosh [25] and Tan and Xu [15] have approximated fixed points of a nonexpansive mapping T by Ishikawa iterates under the condition (A).

In the sequel, the following lemmas are needed to prove our main results.

Lemma 1.5 (see [15]). *Let $\{a_n\}$, $\{b_n\}$ and $\{\delta_n\}$ be sequences of non-negative real numbers satisfying the inequality*

$$a_{n+1} \leq (1 + \delta_n)a_n + b_n, \quad n \geq 1.$$

If $\sum_{n=1}^{\infty} b_n < \infty$ and $\sum_{n=1}^{\infty} \delta_n < \infty$, then

(i) $\lim_{n \rightarrow \infty} a_n$ exists;

(ii) In particular, if $\{a_n\}$ has a sequence $\{a_{n_k}\}$ converging to 0, then

$\lim_{n \rightarrow \infty} a_n = 0$.

Lemma 1.6 (see [12]). *Let X be a real uniformly convex Banach space and $0 \leq p \leq t_n \leq q < 1$ for all positive integer $n \geq 1$. Also suppose that $\{x_n\}$ and $\{y_n\}$ are two sequences of X such that $\limsup_{n \rightarrow \infty} \|x_n\| \leq r$, $\limsup_{n \rightarrow \infty} \|y_n\| \leq r$ and $\lim_{n \rightarrow \infty} \|t_n x_n + (1 - t_n)y_n\| = r$ hold for some $r \geq 0$, then $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$.*

2 Main Results

In order to prove our main results, the following lemmas are needed.

Lemma 2.1. *Let X be a real Banach space and C a nonempty closed convex nonexpansive retract of X with P as a nonexpansive retraction. Let $T_1, T_2 : C \rightarrow X$ be two nonself generalized asymptotically quasi-nonexpansive mappings of C with sequences $\{k_n^{(i)}\}$, $\{\delta_n^{(i)}\} \subset [1, \infty)$ ($i = 1, 2$), respectively such that $\sum_{n=1}^{\infty} (k_n^{(i)} - 1) < \infty$, $\sum_{n=1}^{\infty} \delta_n^{(i)} < \infty$ and $F = F(T_1) \cap F(T_2) \neq \emptyset$. Suppose that $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$ and $\{\lambda_n\}$ are real sequences in $[0, 1)$ such that $\sum_{n=1}^{\infty} \gamma_n < \infty$, $\sum_{n=1}^{\infty} \lambda_n < \infty$ and $\{u_n\}, \{v_n\}$ are bounded sequences in C . From an arbitrary $x_1 \in C$, define the sequence $\{x_n\}$ by (1.17). If $q \in F$, then $\lim_{n \rightarrow \infty} \|x_n - q\|$ exists.*

Proof. Let $q \in F$, by boundedness of the sequences $\{u_n\}$ and $\{v_n\}$, so we can put

$$M = \max\{\sup_{n \geq 1} \|u_n - q\|, \sup_{n \geq 1} \|v_n - q\|\}.$$

Setting $k_n^{(1)} = 1 + r_n^{(1)}$, $k_n^{(2)} = 1 + r_n^{(2)}$. Since $\sum_{n=1}^{\infty} (k_n^{(i)} - 1) < \infty$ ($i = 1, 2$), so $\sum_{n=1}^{\infty} r_n^{(1)} < \infty$, $\sum_{n=1}^{\infty} r_n^{(2)} < \infty$. Using (1.17), we have

$$\begin{aligned}
 \|y_n - q\| &= \|P((1 - \beta_n - \gamma_n)x_n + \beta_n T_2 (PT_2)^{n-1} x_n + \gamma_n v_n) - P(q)\| \\
 &\leq \|(1 - \beta_n - \gamma_n)(x_n - q) + \beta_n (T_2 (PT_2)^{n-1} x_n - q) + \gamma_n (v_n - q)\| \\
 &\leq (1 - \beta_n - \gamma_n)\|x_n - q\| + \beta_n \|T_2 (PT_2)^{n-1} x_n - q\| + \gamma_n \|v_n - q\| \\
 &\leq (1 - \beta_n - \gamma_n)\|x_n - q\| + \beta_n (1 + r_n^{(2)})\|x_n - q\| + \delta_n^{(2)} + \gamma_n M \\
 &= (1 - \beta_n - \gamma_n)\|x_n - q\| + (\beta_n + \beta_n r_n^{(2)})\|x_n - q\| + \delta_n^{(2)} + \gamma_n M \\
 &\leq \|x_n - q\| + r_n^{(2)}\|x_n - q\| + \delta_n^{(2)} + \gamma_n M \\
 &= (1 + r_n^{(2)})\|x_n - q\| + \delta_n^{(2)} + \gamma_n M,
 \end{aligned}$$

and so

$$\begin{aligned}
 \|x_{n+1} - q\| &= \|P((1 - \alpha_n - \lambda_n)y_n + \alpha_n T_1 (PT_1)^{n-1} y_n + \lambda_n u_n) - P(q)\| \\
 &\leq \|(1 - \alpha_n - \lambda_n)(y_n - q) + \alpha_n (T_1 (PT_1)^{n-1} y_n - q) + \lambda_n (u_n - q)\| \\
 &\leq (1 - \alpha_n - \lambda_n)\|y_n - q\| + \alpha_n \|T_1 (PT_1)^{n-1} y_n - q\| + \lambda_n \|u_n - q\| \\
 &\leq (1 - \alpha_n - \lambda_n)\|y_n - q\| + \alpha_n (1 + r_n^{(1)})\|y_n - q\| + \delta_n^{(1)} + \lambda_n M \\
 &= (1 - \alpha_n - \lambda_n)\|y_n - q\| + (\alpha_n + \alpha_n r_n^{(1)})\|y_n - q\| + \delta_n^{(1)} + \lambda_n M \\
 &\leq \|y_n - q\| + r_n^{(1)}\|y_n - q\| + \delta_n^{(1)} + \lambda_n M \\
 &= (1 + r_n^{(1)})\|y_n - q\| + \delta_n^{(1)} + \lambda_n M \\
 &\leq (1 + r_n^{(1)})(1 + r_n^{(2)})\|x_n - q\| + \delta_n^{(2)} + \gamma_n M + \delta_n^{(1)} + \lambda_n M \\
 &= (1 + r_n^{(1)})(1 + r_n^{(2)})\|x_n - q\| \\
 &\quad + (1 + r_n^{(1)})\delta_n^{(2)} + (1 + r_n^{(1)})\gamma_n M + \delta_n^{(1)} + \lambda_n M \\
 &= (1 + r_n^{(1)} + r_n^{(2)} + r_n^{(1)}r_n^{(2)})\|x_n - q\| + \varepsilon_n^{(1)},
 \end{aligned}$$

where $\varepsilon_n^{(1)} = (1 + r_n^{(1)})\delta_n^{(2)} + (1 + r_n^{(1)})\gamma_n M + \delta_n^{(1)} + \lambda_n M$ and we note here that $\sum_{n=1}^\infty \varepsilon_n^{(1)} < \infty$ since $\sum_{n=1}^\infty \gamma_n < \infty$, $\sum_{n=1}^\infty \lambda_n < \infty$, $\sum_{n=1}^\infty r_n^{(1)} < \infty$, $\sum_{n=1}^\infty \delta_n^{(1)} < \infty$ and $\sum_{n=1}^\infty \delta_n^{(2)} < \infty$. Since $\sum_{n=1}^\infty (r_n^{(1)} + r_n^{(2)} + r_n^{(1)}r_n^{(2)}) < \infty$ we obtained by Lemma 1.5 (i) that $\lim_{n \rightarrow \infty} \|x_n - q\|$ exists. This completes the proof. \square

Lemma 2.2. *Let X be a real uniformly convex Banach space and C a nonempty closed convex nonexpansive retract of X with P as a nonexpansive retraction. Let $T_1, T_2 : C \rightarrow X$ be two uniformly L -Lipschitzian, nonself generalized asymptotically quasi-nonexpansive mappings of C with sequences $\{k_n^{(i)}\}, \{\delta_n^{(i)}\} \subset [1, \infty)$ ($i = 1, 2$), respectively such that $\sum_{n=1}^\infty (k_n^{(i)} - 1) < \infty$, $\sum_{n=1}^\infty \delta_n^{(i)} < \infty$ and $F = F(T_1) \cap F(T_2) \neq \emptyset$. Suppose that $\{\alpha_n\}, \{\beta_n\}$ are real sequences in $[\epsilon, 1 - \epsilon]$ for some $\epsilon \in (0, 1)$, $\{\gamma_n\}, \{\lambda_n\} \subset [0, 1)$ such that $\sum_{n=1}^\infty \gamma_n < \infty$, $\sum_{n=1}^\infty \lambda_n < \infty$ and $\{u_n\}, \{v_n\}$ are bounded sequences in C . From an arbitrary $x_1 \in C$, define the sequence $\{x_n\}$ by (1.17). Then $\lim_{n \rightarrow \infty} \|x_n - T_1 x_n\| = \lim_{n \rightarrow \infty} \|x_n - T_2 x_n\| = 0$.*

Proof. Let $q \in F$. Setting $k_n^{(1)} = 1 + r_n^{(1)}$, $k_n^{(2)} = 1 + r_n^{(2)}$. By Lemma 2.1, we see that $\lim_{n \rightarrow \infty} \|x_n - q\|$ exists. It follows that $\{x_n\}$ and $\{y_n\}$ are bounded. Also, $\{u_n - y_n\}$ and $\{v_n - x_n\}$ are bounded. Now we set

$$C = \max\{\sup_{n \geq 1} \|u_n - y_n\|, \sup_{n \geq 1} \|v_n - x_n\|\}.$$

Assume that $\lim_{n \rightarrow \infty} \|x_n - q\| = c$. In addition,

$$\|y_n - q\| \leq (1 + r_n^{(2)})\|x_n - q\| + \delta_n^{(2)} + \gamma_n M, \tag{2.1}$$

where the notation M is taken from Lemma 2.1.

Taking the lim sup on both sides in the inequality (2.1), we have

$$\limsup_{n \rightarrow \infty} \|y_n - q\| \leq c. \tag{2.2}$$

Note that $\|y_n - q + \lambda_n(u_n - y_n)\| \leq \|y_n - q\| + \lambda_n C$ gives that

$$\limsup_{n \rightarrow \infty} \|y_n - q + \lambda_n(u_n - y_n)\| \leq c. \tag{2.3}$$

In addition, $\|T_1(PT_1)^{n-1}y_n - q + \lambda_n(u_n - y_n)\| \leq k_n^{(1)}\|y_n - q\| + \delta_n^{(1)} + \lambda_n C$, taking the lim sup on both sides in this inequality, we have

$$\limsup_{n \rightarrow \infty} \|T_1(PT_1)^{n-1}y_n - q + \lambda_n(u_n - y_n)\| \leq c. \tag{2.4}$$

In addition,

$$\begin{aligned} \|x_{n+1} - q\| &\leq \|(1 - \alpha_n - \lambda_n)(y_n - q) + \alpha_n(T_1(PT_1)^{n-1}y_n - q) + \lambda_n(u_n - q)\| \\ &\leq (1 + r_n^{(1)} + r_n^{(2)} + r_n^{(1)}r_n^{(2)})\|x_n - q\| + \varepsilon_n^{(1)}, \end{aligned} \tag{2.5}$$

where the notation $\varepsilon_n^{(1)}$ is taken from Lemma 2.1.

Since $\sum_{n=1}^{\infty} (r_n^{(1)} + r_n^{(2)} + r_n^{(1)}r_n^{(2)}) < \infty$, $\sum_{n=1}^{\infty} \varepsilon_n^{(1)} < \infty$ and $\lim_{n \rightarrow \infty} \|x_{n+1} - q\| = c$, letting $n \rightarrow \infty$ in the inequality (2.5), we have

$$\lim_{n \rightarrow \infty} \|(1 - \alpha_n - \lambda_n)(y_n - q) + \alpha_n(T_1(PT_1)^{n-1}y_n - q) + \lambda_n(u_n - q)\| = c. \tag{2.6}$$

From

$$\begin{aligned} \|(1 - \alpha_n)(y_n - q + \lambda_n(u_n - y_n)) + \alpha_n(T_1(PT_1)^{n-1}y_n - q + \lambda_n(u_n - y_n))\| = \\ \|(1 - \alpha_n - \lambda_n)(y_n - q) + \alpha_n(T_1(PT_1)^{n-1}y_n - q) + \lambda_n(u_n - q)\|. \end{aligned}$$

and (2.6), we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|(1 - \alpha_n)(y_n - q + \lambda_n(u_n - y_n)) + \alpha_n(T_1(PT_1)^{n-1}y_n - q + \lambda_n(u_n - y_n))\| \\ = c. \end{aligned} \tag{2.7}$$

By using (2.3), (2.4), (2.7) and Lemma 1.6, we have

$$\lim_{n \rightarrow \infty} \|T_1(PT_1)^{n-1}y_n - y_n\| = 0. \quad (2.8)$$

In addition,

$$\begin{aligned} \|T_2(PT_2)^{n-1}x_n - q + \gamma_n(v_n - x_n)\| &\leq \|T_2(PT_2)^{n-1}x_n - q\| + \gamma_n\|v_n - x_n\| \\ &\leq k_n^{(2)}\|x_n - q\| + \delta_n^{(2)} + \gamma_n C, \end{aligned}$$

and taking the lim sup on both sides in this inequality, we have

$$\limsup_{n \rightarrow \infty} \|T_2(PT_2)^{n-1}x_n - q + \gamma_n(v_n - x_n)\| \leq c. \quad (2.9)$$

Using (1.17), we have

$$\begin{aligned} \|x_{n+1} - q\| &\leq (1 - \alpha_n - \lambda_n)\|y_n - q\| + \alpha_n\|T_1(PT_1)^{n-1}y_n - q\| + \lambda_n\|u_n - q\| \\ &= (1 - \alpha_n - \lambda_n)\|y_n - q\| + \alpha_n\|T_1(PT_1)^{n-1}y_n - y_n + y_n - q\| \\ &\quad + \lambda_n\|u_n - y_n + y_n - q\| \\ &\leq (1 - \alpha_n - \lambda_n)\|y_n - q\| + \alpha_n\|T_1(PT_1)^{n-1}y_n - y_n\| \\ &\quad + \alpha_n\|y_n - q\| + \lambda_n\|u_n - y_n\| + \lambda_n\|y_n - q\| \\ &\leq \|y_n - q\| + \|T_1(PT_1)^{n-1}y_n - y_n\| + \lambda_n C. \end{aligned} \quad (2.10)$$

Taking the lim inf on both sides in the inequality (2.10), by (2.8), $\sum_{n=1}^{\infty} \lambda_n < \infty$ and $\lim_{n \rightarrow \infty} \|x_{n+1} - q\| = c$, we have

$$\liminf_{n \rightarrow \infty} \|y_n - q\| \geq c. \quad (2.11)$$

It follows from (2.2) and (2.11) that $\lim_{n \rightarrow \infty} \|y_n - q\| = c$. This implies that

$$\begin{aligned} c = \lim_{n \rightarrow \infty} \|y_n - q\| &\leq \lim_{n \rightarrow \infty} \|(1 - \beta_n - \gamma_n)(x_n - q) \\ &\quad + \beta_n(T_2(PT_2)^{n-1}x_n - q) + \gamma_n(v_n - q)\| \\ &\leq \lim_{n \rightarrow \infty} \|x_n - q\| = c, \end{aligned}$$

and so

$$\lim_{n \rightarrow \infty} \|(1 - \beta_n - \gamma_n)(x_n - q) + \beta_n(T_2(PT_2)^{n-1}x_n - q) + \gamma_n(v_n - q)\| = c. \quad (2.12)$$

From

$$\begin{aligned} \|(1 - \beta_n)(x_n - q + \gamma_n(v_n - x_n)) + \beta_n(T_2(PT_2)^{n-1}x_n - q + \gamma_n(v_n - x_n))\| \\ = \|(1 - \beta_n - \gamma_n)(x_n - q) + \beta_n(T_2(PT_2)^{n-1}x_n - q) + \gamma_n(v_n - q)\| \end{aligned}$$

and (2.12), we have

$$\lim_{n \rightarrow \infty} \|(1 - \beta_n)(x_n - q + \gamma_n(v_n - x_n)) + \beta_n(T_2(PT_2)^{n-1}x_n - q + \gamma_n(v_n - x_n))\| = c. \quad (2.13)$$

Note that $\|x_n - q + \gamma_n(v_n - x_n)\| \leq \|x_n - q\| + \gamma_n C$ gives that

$$\limsup_{n \rightarrow \infty} \|x_n - q + \gamma_n(v_n - x_n)\| \leq c. \tag{2.14}$$

Using (2.9), (2.13), (2.14) and Lemma 1.6, we obtain

$$\lim_{n \rightarrow \infty} \|T_2(PT_2)^{n-1}x_n - x_n\| = 0. \tag{2.15}$$

From $y_n = P((1 - \beta_n - \gamma_n)x_n + \beta_n T_2(PT_2)^{n-1}x_n + \gamma_n v_n)$, $\sum_{n=1}^{\infty} \gamma_n < \infty$ and (2.15), we have

$$\begin{aligned} \|y_n - x_n\| &= \|P((1 - \beta_n - \gamma_n)x_n + \beta_n T_2(PT_2)^{n-1}x_n + \gamma_n v_n) - x_n\| \\ &\leq \|(1 - \beta_n - \gamma_n)(x_n - x_n) + \beta_n(T_2(PT_2)^{n-1}x_n - x_n) + \gamma_n(v_n - x_n)\| \\ &\leq \beta_n \|T_2(PT_2)^{n-1}x_n - x_n\| + \gamma_n \|v_n - x_n\| \\ &\leq \|T_2(PT_2)^{n-1}x_n - x_n\| + \gamma_n C \\ &\rightarrow 0 \quad (\text{as } n \rightarrow \infty). \end{aligned} \tag{2.16}$$

Now, since $T_i (i = 1, 2)$ are uniformly L -Lipschitzian for Lipschitz constant $L = \max_{1 \leq i \leq 2} \{L_i\} > 0$. We note that

$$\begin{aligned} \|T_1(PT_1)^{n-1}x_n - x_n\| &= \|T_1(PT_1)^{n-1}x_n - y_n + y_n - x_n\| \\ &\leq \|T_1(PT_1)^{n-1}x_n - y_n\| + \|y_n - x_n\| \\ &= \|T_1(PT_1)^{n-1}x_n - T_1(PT_1)^{n-1}y_n \\ &\quad + T_1(PT_1)^{n-1}y_n - y_n\| + \|y_n - x_n\| \\ &\leq \|T_1(PT_1)^{n-1}x_n - T_1(PT_1)^{n-1}y_n\| \\ &\quad + \|T_1(PT_1)^{n-1}y_n - y_n\| + \|y_n - x_n\| \\ &\leq L\|x_n - y_n\| + \|T_1(PT_1)^{n-1}y_n - y_n\| + \|y_n - x_n\|. \end{aligned}$$

Thus, it follows from (2.8) and (2.16) that

$$\lim_{n \rightarrow \infty} \|T_1(PT_1)^{n-1}x_n - x_n\| = 0. \tag{2.17}$$

By using (1.17), we have

$$\begin{aligned} \|x_{n+1} - x_n\| &\leq (1 - \alpha_n - \lambda_n)\|y_n - x_n\| + \alpha_n \|T_1(PT_1)^{n-1}y_n - x_n\| + \lambda_n \|u_n - x_n\| \\ &\leq (1 - \alpha_n - \lambda_n)\|y_n - x_n\| + \alpha_n \|T_1(PT_1)^{n-1}y_n - y_n + y_n - x_n\| \\ &\quad + \lambda_n \|u_n - y_n + y_n - x_n\| \\ &\leq (1 - \alpha_n - \lambda_n)\|y_n - x_n\| + \alpha_n \|T_1(PT_1)^{n-1}y_n - y_n\| \\ &\quad + \alpha_n \|y_n - x_n\| + \lambda_n \|u_n - y_n\| + \lambda_n \|y_n - x_n\| \\ &\leq \|y_n - x_n\| + \|T_1(PT_1)^{n-1}y_n - y_n\| + \lambda_n C. \end{aligned}$$

It follows from (2.8), (2.16) and $\sum_{n=1}^{\infty} \lambda_n < \infty$ that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \quad (2.18)$$

Using (2.17) and (2.18), we have

$$\begin{aligned} \|x_{n+1} - T_1(PT_1)^{n-1}x_{n+1}\| &= \|x_{n+1} - x_n + x_n - T_1(PT_1)^{n-1}x_n \\ &\quad + T_1(PT_1)^{n-1}x_n - T_1(PT_1)^{n-1}x_{n+1}\| \\ &\leq \|x_{n+1} - x_n\| + \|T_1(PT_1)^{n-1}x_{n+1} - T_1(PT_1)^{n-1}x_n\| \\ &\quad + \|T_1(PT_1)^{n-1}x_n - x_n\| \\ &\leq \|x_{n+1} - x_n\| + L\|x_{n+1} - x_n\| + \|T_1(PT_1)^{n-1}x_n - x_n\|, \\ &\rightarrow 0 \quad (\text{as } n \rightarrow \infty). \end{aligned} \quad (2.19)$$

In addition, for $n \geq 2$,

$$\begin{aligned} \|x_{n+1} - T_1(PT_1)^{n-2}x_{n+1}\| &= \|x_{n+1} - x_n + x_n - T_1(PT_1)^{n-2}x_n \\ &\quad + T_1(PT_1)^{n-2}x_n - T_1(PT_1)^{n-2}x_{n+1}\| \\ &\leq \|x_{n+1} - x_n\| + \|T_1(PT_1)^{n-2}x_n - x_n\| \\ &\quad + \|T_1(PT_1)^{n-2}x_{n+1} - T_1(PT_1)^{n-2}x_n\| \\ &\leq \|x_{n+1} - x_n\| + \|T_1(PT_1)^{n-2}x_n - x_n\| \\ &\quad + L\|x_{n+1} - x_n\|. \end{aligned}$$

It follows from (2.18) and (2.19) that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - T_1(PT_1)^{n-2}x_{n+1}\| = 0. \quad (2.20)$$

We denote as $(PT_1)^{1-1}$ the identity maps from C onto itself. Thus by the inequality (2.19) and (2.20), we have

$$\begin{aligned} \|x_{n+1} - T_1x_{n+1}\| &= \|x_{n+1} - T_1(PT_1)^{n-1}x_{n+1} + T_1(PT_1)^{n-1}x_{n+1} - T_1x_{n+1}\| \\ &\leq \|x_{n+1} - T_1(PT_1)^{n-1}x_{n+1}\| + \|T_1(PT_1)^{n-1}x_{n+1} - T_1x_{n+1}\| \\ &\leq \|x_{n+1} - T_1(PT_1)^{n-1}x_{n+1}\| + L\|(PT_1)^{n-1}x_{n+1} - x_{n+1}\| \\ &= \|x_{n+1} - T_1(PT_1)^{n-1}x_{n+1}\| + L\|(PT_1)(PT_1)^{n-2}x_{n+1} - P(x_{n+1})\| \\ &\leq \|x_{n+1} - T_1(PT_1)^{n-1}x_{n+1}\| + L\|T_1(PT_1)^{n-2}x_{n+1} - x_{n+1}\| \\ &\rightarrow 0 \quad (\text{as } n \rightarrow \infty), \end{aligned}$$

which implies that $\lim_{n \rightarrow \infty} \|x_n - T_1x_n\| = 0$. Similarly, we may show that

$$\lim_{n \rightarrow \infty} \|x_n - T_2x_n\| = 0.$$

The proof is completed. □

We prove the strong convergence of the scheme (1.17) under condition A' which is weaker than the compactness of the domain of the mappings.

Theorem 2.3. *Let X be a real uniformly convex Banach space and C a nonempty closed convex nonexpansive retract of X with P as a nonexpansive retraction. Let $T_1, T_2 : C \rightarrow X$ be two uniformly L -Lipschitzian, nonself generalized asymptotically quasi-nonexpansive mappings of C satisfying condition A' with sequences $\{k_n^{(i)}\}, \{\delta_n^{(i)}\} \subset [1, \infty)$ ($i = 1, 2$), respectively such that $\sum_{n=1}^{\infty} (k_n^{(i)} - 1) < \infty$, $\sum_{n=1}^{\infty} \delta_n^{(i)} < \infty$ and $F = F(T_1) \cap F(T_2) \neq \emptyset$. Suppose that $\{\alpha_n\}, \{\beta_n\}$ are real sequences in $[\epsilon, 1 - \epsilon]$ for some $\epsilon \in (0, 1)$, $\{\gamma_n\}, \{\lambda_n\} \subset [0, 1)$ such that $\sum_{n=1}^{\infty} \gamma_n < \infty$, $\sum_{n=1}^{\infty} \lambda_n < \infty$ and $\{u_n\}, \{v_n\}$ are bounded sequences in C . Then the sequences $\{x_n\}$ and $\{y_n\}$ defined by the iterative scheme (1.17) converge strongly to a common fixed point of T_1 and T_2 .*

Proof. By Lemma 2.2, we have $\lim_{n \rightarrow \infty} \|x_n - T_1 x_n\| = \lim_{n \rightarrow \infty} \|x_n - T_2 x_n\| = 0$. It follows from condition A' that

$$\begin{aligned} \lim_{n \rightarrow \infty} f(d(x_n, F)) &\leq \lim_{n \rightarrow \infty} \|x_n - T_1 x_n\| = 0 \quad \text{or} \\ \lim_{n \rightarrow \infty} f(d(x_n, F)) &\leq \lim_{n \rightarrow \infty} \|x_n - T_2 x_n\| = 0. \end{aligned}$$

In the both case, $\lim_{n \rightarrow \infty} f(d(x_n, F)) = 0$. Since $f : [0, \infty) \rightarrow [0, \infty)$ is a nondecreasing function satisfying $f(0) = 0$, $f(r) > 0$ for all $r \in (0, \infty)$, we obtain that $\lim_{n \rightarrow \infty} d(x_n, F) = 0$. That is

$$\lim_{n \rightarrow \infty} \inf_{y^* \in F} \|x_n - y^*\| = \lim_{n \rightarrow \infty} d(x_n, F) = 0.$$

It implies that

$$\inf_{y^* \in F} \lim_{n \rightarrow \infty} \|x_n - y^*\| = 0.$$

So, for any given $\epsilon > 0$, there exists $p \in F$ and $N > 0$ such that for all $n \geq N$ $\|x_n - p\| < \frac{\epsilon}{2}$. This shows that

$$\begin{aligned} \|x_{n+m} - x_n\| &\leq \|x_{n+m} - p\| + \|x_n - p\| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \end{aligned}$$

for all $n \geq N$ and $m \geq 1$. Hence, $\{x_n\}$ is a Cauchy sequence and so is convergent since X is complete. Let $\lim_{n \rightarrow \infty} x_n = u$. From $\lim_{n \rightarrow \infty} \|x_n - T_1 x_n\| = \lim_{n \rightarrow \infty} \|x_n - T_2 x_n\| = 0$ and the continuity of T_1 and T_2 , we have $\|T_1 u - u\| = \|T_2 u - u\| = 0$. Thus $u \in F$. From (2.16), we have

$$\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0,$$

it follows that $\lim_{n \rightarrow \infty} \|y_n - u\| = 0$. This completes the proof. □

The following result follows from Theorem 2.3.

Theorem 2.4. *Let X be a real uniformly convex Banach space and C a nonempty closed convex nonexpansive retract of X with P as a nonexpansive retraction. Let $T_1, T_2 : C \rightarrow X$ be two nonself asymptotically nonexpansive mappings of C satisfying condition A' with sequences $\{k_n^{(i)}\} \subset [1, \infty)$ ($i = 1, 2$) such that $\sum_{n=1}^{\infty} (k_n^{(i)} - 1) < \infty$ and $F = F(T_1) \cap F(T_2) \neq \emptyset$. Suppose that $\{\alpha_n\}, \{\beta_n\}$ are real sequences in $[\epsilon, 1 - \epsilon]$ for some $\epsilon \in (0, 1)$, $\{\gamma_n\}, \{\lambda_n\} \subset [0, 1)$ such that $\sum_{n=1}^{\infty} \gamma_n < \infty$, $\sum_{n=1}^{\infty} \lambda_n < \infty$ and $\{u_n\}, \{v_n\}$ are bounded sequences in C . Then the sequences $\{x_n\}$ and $\{y_n\}$ defined by the iterative scheme (1.17) converge strongly to a common fixed point of T_1 and T_2 .*

For $\gamma_n = \lambda_n = 0$, the iterative scheme (1.17) reduces to that of (1.16) for uniformly L -Lipschitzian, nonself generalized asymptotically quasi-nonexpansive mappings and the following result is directly obtained by Theorem 2.3.

Theorem 2.5. *Let X be a real uniformly convex Banach space and C a nonempty closed convex nonexpansive retract of X with P as a nonexpansive retraction. Let $T_1, T_2 : C \rightarrow X$ be two uniformly L -Lipschitzian, nonself generalized asymptotically quasi-nonexpansive mappings of C satisfying condition A' with sequences $\{k_n^{(i)}\}, \{\delta_n^{(i)}\} \subset [1, \infty)$ ($i = 1, 2$), respectively such that $\sum_{n=1}^{\infty} (k_n^{(i)} - 1) < \infty$, $\sum_{n=1}^{\infty} \delta_n^{(i)} < \infty$ and $F = F(T_1) \cap F(T_2) \neq \emptyset$. Suppose that $\{\alpha_n\}$ and $\{\beta_n\}$ are real sequences in $[\epsilon, 1 - \epsilon]$ for some $\epsilon \in (0, 1)$. Then the sequences $\{x_n\}$ and $\{y_n\}$ defined by the iterative scheme (1.16) converge strongly to a common fixed point of T_1 and T_2 .*

The following result follows from Theorem 2.5.

Theorem 2.6. *Let X be a real uniformly convex Banach space and C a nonempty closed convex nonexpansive retract of X with P as a nonexpansive retraction. Let $T_1, T_2 : C \rightarrow X$ be two nonself asymptotically nonexpansive mappings of C satisfying condition A' with sequences $\{k_n^{(i)}\} \subset [1, \infty)$ ($i = 1, 2$) such that $\sum_{n=1}^{\infty} (k_n^{(i)} - 1) < \infty$ and $F = F(T_1) \cap F(T_2) \neq \emptyset$. Suppose that $\{\alpha_n\}$ and $\{\beta_n\}$ are real sequences in $[\epsilon, 1 - \epsilon]$ for some $\epsilon \in (0, 1)$. Then the sequences $\{x_n\}$ and $\{y_n\}$ defined by the iterative scheme (1.16) converge strongly to a common fixed point of T_1 and T_2 .*

In the remainder of this section, we deal with the strong convergence of the new iterative scheme (1.17) to a common fixed point of nonself generalized asymptotically quasi-nonexpansive mappings in a real Banach space.

Theorem 2.7. *Let X be a real Banach space and C a nonempty closed convex nonexpansive retract of X with P as a nonexpansive retraction. Let $T_1, T_2 : C \rightarrow X$ be two nonself generalized asymptotically quasi-nonexpansive mappings of C with sequences $\{k_n^{(i)}\}, \{\delta_n^{(i)}\} \subset [1, \infty)$ ($i = 1, 2$), respectively such that $\sum_{n=1}^{\infty} (k_n^{(i)} - 1) < \infty$, $\sum_{n=1}^{\infty} \delta_n^{(i)} < \infty$ and $F = F(T_1) \cap F(T_2) \neq \emptyset$ is closed. Suppose that $\{\alpha_n\}, \{\beta_n\}$ are real sequences in $[\epsilon, 1 - \epsilon]$ for some $\epsilon \in (0, 1)$, $\{\gamma_n\}, \{\lambda_n\} \subset [0, 1)$ such that $\sum_{n=1}^{\infty} \gamma_n < \infty$, $\sum_{n=1}^{\infty} \lambda_n < \infty$ and $\{u_n\}, \{v_n\}$ are bounded sequences in C . Then*

the sequence $\{x_n\}$ defined by the iterative scheme (1.17) converges strongly to a common fixed point of T_1 and T_2 if and only if $\liminf_{n \rightarrow \infty} d(x_n, F) = 0$, where $d(x_n, F) = \inf_{y \in F} \|x_n - y\|$, $n \geq 1$.

Proof. The necessity of the conditions is obvious. Thus, we will only prove the sufficiency. As in the proof of Lemma 2.1, by the arbitrariness of $q \in F$, we have

$$\|x_{n+1} - q\| \leq (1 + r_n^{(1)} + r_n^{(2)} + r_n^{(1)}r_n^{(2)})\|x_n - q\| + \varepsilon_n^{(1)},$$

and so

$$d(x_{n+1}, F) \leq (1 + r_n^{(1)} + r_n^{(2)} + r_n^{(1)}r_n^{(2)})d(x_n, F) + \varepsilon_n^{(1)},$$

where $\varepsilon_n^{(1)} = (1 + r_n^{(1)})\delta_n^{(2)} + (1 + r_n^{(1)})\gamma_n M + \delta_n^{(1)} + \lambda_n M$. Since $\sum_{n=1}^{\infty} (r_n^{(1)} + r_n^{(2)} + r_n^{(1)}r_n^{(2)}) < \infty$ and $\sum_{n=1}^{\infty} \varepsilon_n^{(1)} < \infty$, we obtained by Lemma 1.5 that $\lim_{n \rightarrow \infty} d(x_n, F)$ exists. Then, by hypothesis $\liminf_{n \rightarrow \infty} d(x_n, F) = 0$, we have $\lim_{n \rightarrow \infty} d(x_n, F) = 0$. From Theorem 2.3, it obtain that $\{x_n\}$ defined by (1.17) is a Cauchy sequence in C . Let $\lim_{n \rightarrow \infty} x_n = u$. Now $\lim_{n \rightarrow \infty} d(x_n, F) = 0$ gives that $d(u, F) = 0$. F is closed; therefore $u \in F$. This completes the proof of Theorem 2.7. \square

Acknowledgement : This research was supported by the revenue budget School of Science, University of Phayao. T. Thianwan would like to thank University of Phayao, Phayao, Thailand (Grant No. UoE62001).

References

- [1] C.E. Chidume, B. Ali, Convergence theorems for finite families of asymptotically quasi-nonexpansive mappings, *Journal of Inequalities and Application* 2007 (2007) DOI: 10.1155/2007/68616.
- [2] C.E. Chidume, E.U. Ofoedu, H. Zegeye, Strong and weak convergence theorems for asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* 280 (2003) 364-374.
- [3] L.B. Ćirić, J.S. Ume, S.N. Ješi, M.M. Arandjelović-Milovanović, *Numer. Funct. Anal. Optim.* 28 (11-12) (2007) 1231-1243.
- [4] L.B. Ćirić, J.S. Ume, *Acta Math. Hungar.* 98 (1-2) (2003) 1-8.
- [5] L.B. Ćirić, J.S. Ume, M.S. Khan, *Archivum Mathematicum (Brno)*. 39 (2003) 123-127.
- [6] G. Das, J.P. Debata, Fixed points of quasi-nonexpansive mappings, *Indian J. Pure. Appl. Math.* 17 (1986) 1263-1269.
- [7] W.J. Davis, P. Enflo, Contractive projections on lp spaces, *Analysis at Urbana, Vol. I (Urbana, IL, 1986-1987)*, 151161, London Math. Soc. Lecture Note Ser., 137, Cambridge Univ. Press, Cambridge, 1989.

- [8] W. Takahashi, T. Tamura, Limit theorems of operators by convex combinations of nonexpansive retractions in Banach space, *J. Approx. Theory.* 91 (3) (1997) 386-397.
- [9] S.H. Khan, W. Takahashi, Approximating common fixed points of two asymptotically nonexpansive mappings, *Sci. Math. Jpn.* 53 (1) (2001) 143-148.
- [10] W. Takahashi, Iterative methods for approximation of fixed points and their applications, *J. Oper. Res. Soc. Jpn.* 43 (1) (2000) 87-108.
- [11] K. Goebel, W.A. Kirk, A fixed point theorem for asymptotically nonexpansive mappings, *Proc. Amer. Math. Soc.* 35 (1972) 171-174.
- [12] J. Schu, Weak and strong convergence to fixed points of asymptotically non-expansive mappings, *Bull. Austral. Math. Soc.* 43 (1991) 53-159.
- [13] M.O. Osilike, S.C. Aniagbosor, Weak and strong convergence theorems for fixed points of asymptotically nonexpansive mappings, *Math. Comput. Modelling.* 32 (2000) 1181-1191.
- [14] B.E. Rhoades, Fixed points iteration for certain nonlinear mapping, *J. Math. Anal. Appl.* 183 (1994) 118-120.
- [15] K.K. Tan, H.K. Xu, Approximation fixed points of nonexpansive mappings by the Ishikawa iteration process, *J. Math. Anal. Appl.* 178 (1993) 301-308.
- [16] N. Shahzad, H. Zegeye, Strong convergence of an implicit iteration process for a finite family of generalized asymptotically quasi-nonexpansive mappings, *Appl. Math. Comput.* 189 (2007) 1058-1065.
- [17] W.R. Mann, Mean value methods in iteration, *Proc. Amer. Math. Soc.* 4 (1953) 506-510.
- [18] S.Y. Matsushita, D. Kuroiwa, Strong convergence of averaging iteration of nonexpansive nonself-mappings, *J. Math. Anal. Appl.* 294 (2004) 206-214.
- [19] J. Schu, Weak and strong convergence to fixed points of asymptotically non-expansive mappings, *Bull. Austral. Math. Soc.* 43 (1991) 53-159.
- [20] N. Shahzad, Approximating fixed point of non-self nonexpansive mappings on Banach space *Nonlinear Anal.* 61 (2005) 1031-1039.
- [21] L. Wang, Strong and weak convergence theorems for common fixed points of nonself asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* 323 (2006) 550-557.
- [22] L. Deng, Q. Liu, Iterative scheme for nonself generalized asymptotically quasi-nonexpansive mappings, *Appl. Math. Comput.* 205 (2008) 317-324.
- [23] S. Thianwan, Common fixed points of new iterations for two asymptotically nonexpansive nonself-mappings in Banach spaces, *J. Comput. Appl. Math.* 224 (2009) 688-695.

- [24] H. Fukhar-ud-din, S.H. Khan, Convergence of iterates with errors of asymptotically quasi-nonexpansive mappings and applications, *J. Math. Anal. Appl.* 328 (2007) 821-829.
- [25] M. Maiti, M.K. Ghosh, Approximating fixed points by Ishikwa iterates, *Bull. Austral. Math. Soc.* 40 (1989) 113-117.

(Received 14 June 2019)

(Accepted 4 October 2019)